



## IPKine™ HRP, Goat Anti-Mouse IgG LCS

Cat #: A25012

Size: 100µl /500µl

### Product Information

	<b>Product Name:</b> IPKine™ HRP, Goat Anti-Mouse IgG LCS		
	<b>Applications:</b> WB, IP		<b>Isotype:</b> Goat IgG
	<b>Reactivity:</b> Mouse		
<b>REF</b>	<b>Catalog Number:</b> A25012	<b>LOT</b>	<b>Lot Number:</b> Refer to product label
	<b>Formulation:</b> Liquid		<b>Concentration:</b> 0.5 mg/ml
	<b>Storage:</b> Store at -20°C. Avoid repeated freeze / thaw cycles.		<b>Note:</b> Contain sodium azide.

**Background:** Abbkineline secondary antibodies are available conjugated to enzyme, biotin or fluorophore for use in a variety of antibody-based applications including Western Blot, ImmunoHistoChemistry, ImmunoFluorescence, Flow Cytometry and ELISA. We offer high quality secondary antibodies from goat, rabbit and donkey sources for your each application.

**Application Notes:** The suggested dilution ratio is: WB: 1:1,000-1:10,000 (1:2000 optimal), IP: 1:1,000-1:10,000 (1:2000 optimal). In rare cases, due to the aggregation of light chains, there are weak bands at 50 kDa. It is recommended to use the following method to reduce the non-specific interference background: Increasing the concentration of target protein in the sample, reducing the use of primary antibody in Pull Down, or optimizing the dilution of WB antibody and ECL exposure conditions.

**Storage Buffer:** Liquid in PBS, pH 7.4, containing 1% BSA as stabilizer and 50% Glycerol.

**Storage Instructions:** Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

**Note:** The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

**Example:** Sample: Mouse IgG, Dilution factor:1:2000, Exposure time: 30 s

