



Anti-Myc Tag Mouse Monoclonal Antibody (2D5), Magnetic Beads

Cat #: A02060MGB

Size: 100ul/400ul/1ml

Product Information

	Product Name: Anti-Myc Tag Mouse Monoclonal Antibody (2D5), Magnetic Beads		
	Applications: IP		Isotype: Mouse IgG1
REF	Catalog Number: A02060MGB	LOT	Lot Number: Refer to product label
	Storage: Store at 4°C. Avoid freeze-thaw or centrifugation.		Note: Contain sodium azide.
	Formulation: Over 1 mg of Antibody coupled to 1 ml of packed Magnetic beads.		

Product Description: Anti-Myc Tag Magnetic Beads are prepared by covalently coupling Anti-Myc Tag Mouse Monoclonal Antibody to crosslinked Magnetic Beads, useful for detection and capture of fusion proteins containing a Myc peptide sequence by commonly used immunoprecipitation procedures. The coupling technique is optimized to give a high binding capacity for Myc-tag protein.

Storage Buffer: Suspended in PBS, pH 7.4, containing 0.02% Sodium Azide as preservative.

Storage instructions: Stable for one year at 4°C from date of shipment. Avoid freeze-thaw or centrifugation.

Shipping: Gel pack with blue ice.

Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

Suggested Procedure

Magnetic Beads Preparation (Repeat for three times)

1. Transfer the appropriate amount of Magnetic beads (Suggested volume: 10-20 μ L Magnetic beads for each sample) into a 1.5 mL microfuge tube and place it on the Magnetic separator for 10s.
2. Remove the storage solution.
3. Resuspend the Magnetic beads with 1 mL ice-cold 1 \times TBS.
4. Place it on the Magnetic separator for 10 s and remove the supernatant.

Note: For multiple samples, it is recommended to prepare the Magnetic beads first, and then aliquot into each microfuge tube.

Sample preparation

1. Collect 2×10^6 cells and wash with PBS for three times.
2. Resuspend cells in 400 μ L ice-cold lysis buffer [50 mM Tris (pH 7.5), 150 mM NaCl, 0.05% NP-40], then sonicate briefly (up to 10s).
3. Centrifugate 12000g for 5 minutes at 4°C and collect the supernatant.

Binding protein

1. Add 400 μ L cell lysate to Magnetic beads.
2. Mix thoroughly and incubate at 4°C for 1-2 h.

Note: During the binding process, the Magnetic beads may be clustered or flake. This phenomenon is normal and will not affect the experimental results.

Wash (Repeat 3-5 times until OD280 of the supernatant is lesser than 0.05)

1. Place the tube on the Magnetic separator for 10s to collect the mixture.
2. Resuspend the mixture with 1 mL ice-cold 1×TBS and incubate it at 4°C for 1-3 minutes.
3. Place it on the Magnetic separator for 10 s and remove the supernatant. (Or save the supernatant for further analysis)

Note: Avoid losing Magnetic Beads during wash step.

Elution

For SDS-PAGE detection -- Add 50 μ L 1×protein sample buffer to the above obtained precipitate, and then boil for 5 min. Cool to room temperature and place in on the Magnetic separator for 10s.

For other assay -- Add 2 volumes (vs beads volume) elution buffer [0.1 M -0.2 M Glycine pH 2.5-3.1 (or 0.1 M citric acid, pH 2.5-3.1 or 2.5% Acetic Acid)] to the above obtained precipitate, and incubate at least 2 minutes to collect elution fraction. (Repeat for three times).