Technical support: support@abbkine.com

Website: https://www.abbkine.com

IKB β Monoclonal Antibody

Cat #: ABM40229 Size: 30µl /100µl /200µl

Product Information

| | Product Name: IκB β Monoclonal Antibody | | |
|-----|--|-----|------------------------------------|
| | Applications: IHC-P, IF | | Isotype: Mouse IgG1 |
| | Reactivity: Human, Mouse, Rat | | |
| REF | Catalog Number: ABM40229 | LOT | Lot Number: Refer to product label |
| | Formulation: Liquid | | Concentration: 1 mg/ml |
| Å. | Storage: Store at -20°C. Avoid repeated freeze / thaw cycles. | A | Note: Contain sodium azide. |

<u>Background</u>: The protein encoded by NFKBIB (NFKB inhibitor beta) belongs to the NF-kappa-B inhibitor family, which inhibit NF-kappa-B by complexing with, and trapping it in the cytoplasm. Phosphorylation of serine residues on these proteins by kinases marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B, which translocates to the nucleus to function as a transcription factor. Alternatively spliced transcript variants have been found for NFKBIB.

<u>Application Notes</u>: Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: IHC-P (1:100-1:200).

Storage Buffer: PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.

Storage Instructions: Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.




Fig.1. Immunofluorescence analysis of human liver cancer tissue. 1, IkB β Monoclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 Labeled secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. P






Fig.2. Immunofluorescence analysis of mouse testis tissue. 1, IkB β Monoclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 Labeled secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B.

<u>Note:</u> The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

