Technical support: support@abbkine.com

Website: https://www.abbkine.com

Collagen IV Mouse Monoclonal Antibody (8E5)

Cat #: ABM40371 Size: 30µl /100µl /200µl

Product Information

	Product Name: Collagen IV Mouse Monoclonal Antibody (8E5)		
	Applications: IHC-P, IF		Isotype: Mouse IgG1
	Reactivity: Human, Mouse, Rat		
REF	Catalog Number: ABM40371	LOT	Lot Number: Refer to product label
	Formulation: Liquid		Concentration: 1 mg/ml
Ĵ.	Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.	A	Note: Contain sodium azide.

Background: COL4A1 (collagen type IV alpha 1 chain) encodes a type IV collagen alpha protein. Type IV collagen proteins are integral components of basement membranes. COL4A1 shares a bidirectional promoter with a paralogous gene on the opposite strand. The protein consists of an amino-terminal 7S domain, a triple-helix forming collagenous domain, and a carboxy-terminal non-collagenous domain. It functions as part of a heterotrimer and interacts with other extracellular matrix components such as perlecans, proteoglycans, and laminins. In addition, proteolytic cleavage of the non-collagenous carboxy-terminal domain results in a biologically active fragment known as arresten, which has anti-angiogenic and tumor suppressor properties. Mutations in COL4A1 cause porencephaly, cerebrovascular disease, and renal and muscular defects. Alternative splicing results in multiple transcript variants.

<u>Application Notes</u>: Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: IHC-P (1:50-1:200).

Storage Buffer: PBS, pH 7.4, containing 0.02% Sodium Azide as preservative and 50% Glycerol.

Storage Instructions: Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.






Fig.1. Immunofluorescence analysis of human liver tissue. 1, Collagen IV Mouse Monoclonal Antibody (8E5) (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 Labeled secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blu




Fig.2. Immunofluorescence analysis of mouse spleen tissue. 1, Collagen IV Mouse Monoclonal Antibody (8E5) (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 Labeled secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B.

Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

