Technical support: support@abbkine.com

Website: https://www.abbkine.com

ATG7 Mouse Monoclonal Antibody (3D6)

Cat #: ABM40378 Size: 30µl /100µl /200µl

Product Information

	Product Name: ATG7 Mouse Monoclonal Antibody (3D6)		
	Applications: IHC-P, IF		Isotype: Mouse IgG1
	Reactivity: Human, Mouse, Rat		
REF	Catalog Number: ABM40378	LOT	Lot Number: Refer to product label
	Formulation: Liquid		Concentration: 1 mg/ml
Å	Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.	<u>∧</u>	Note: Contain sodium azide.

Background: ATG7 (autophagy related 7) encodes an E1-like activating enzyme that is essential for autophagy and cytoplasmic to vacuole transport. The encoded protein is also thought to modulate p53-dependent cell cycle pathways during prolonged metabolic stress. It has been associated with multiple functions, including axon membrane trafficking, axonal homeostasis, mitophagy, adipose differentiation, and hematopoietic stem cell maintenance. Alternative splicing results in multiple transcript variants.

<u>Application Notes</u>: Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: IHC-P (1:50-1:200).

Storage Buffer: PBS, pH 7.4, containing 0.02% Sodium Azide as preservative and 50% Glycerol.

Storage Instructions: Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.




Fig.1. Immunofluorescence analysis of mouse spleen tissue. 1, ATG7 Mouse Monoclonal Antibody (3D6) (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 Labeled secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10m






Fig.2. Immunohistochemical analysis of paraffinembedded human uterus tissue. 1, ATG7 Mouse Monoclonal Antibody (3D6) was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.

Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

