Customer service: service@abbkine.com
Technical support: support@abbkine.com
Website: https://www.abbkine.com

AMPKα1/2 Polyclonal Antibody

Cat #: ABP0060 Size: 100μl

Product Information

	Product Name: AMPKα1/2 Polyclonal Antibody		
	Applications: WB, IHC-P, ELISA		**Isotype:** Rabbit IgG
	Reactivity: Human, Mouse, Rat, Monkey		
REF	**Catalog Number:** ABP0060	LOT	**Lot Number:** Refer to product label
	Formulation: Liquid		**Concentration:** 1 mg/ml
	Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.		**Note:** Contain sodium azide.

Background: Protein kinase AMP-activated catalytic subunit alpha 1 encoded by PRKAA1 belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all eukaryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio. AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protects cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alternatively spliced transcript variants encoding distinct isoforms have been observed.

Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: WB (1:500-1:2000), IHC-P (1:100-1:300), ELISA (1:40000). Not yet tested in other applications.

Storage Buffer: PBS, pH 7.4, containing 0.02% Sodium Azide as preservative and 50% Glycerol.

Storage Instructions: Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

Not for further distribution without written consent.Copyright © Abbkine, Inc.

	Fig.1. Western blot analysis of various cells using AMPKα1/2 Polyclonal Antibody diluted at 1:500.
	Fig.2. Immunohistochemical analysis of paraffin-embedded mouse testis tissue. 1, AMPKα1/2 Polyclonal Antibody was diluted at 1:200 (4°C, overnight). 2, Sodium citrate pH 6.0 was used for antibody retrieval (>98°C, 20min). 3, secondary antibody was diluted at 1:200 (room temperature, 30min). Negative control was used by secondary antibody only.

Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

Abbkine, Inc • China • ***www.abbkine.com***
Not for further distribution without written consent.Copyright © Abbkine, Inc.