Customer service: service@abbkine.com
Technical support: support@abbkine.com
Website: https://www.abbkine.com

Smad2 Polyclonal Antibody

Cat #: ABP0158 Size: 100μl

Product Information

	Product Name: Smad2 Polyclonal Antibody		
	Applications: WB, IF, IHC-P, ELISA		**Isotype:** Rabbit IgG
	Reactivity: Human, Mouse, Rat		
REF	**Catalog Number:** ABP0158	LOT	**Lot Number:** Refer to product label
	Formulation: Liquid		**Concentration:** 1 mg/ml
	Storage: Store at -20°C. Avoid repeated freeze / thaw cycles.		**Note:** Contain sodium azide.

Background: The protein encoded by SMAD2 belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene ‘mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene.

Application Notes: Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: WB (1:500-1:2000), IF (1:200-1:1000), IHC-P (1:100-1:300), ELISA (1:5000). Not yet tested in other applications.

Storage Buffer: PBS, pH 7.4, containing 0.02% Sodium Azide as preservative and 50% Glycerol.

Storage Instructions: Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

Not for further distribution without written consent.Copyright © Abbkine, Inc.

	Fig.1. Western blot analysis of COLO cells using Smad2 Polyclonal Antibody diluted at 1:1000.
	Fig.2. Immunofluorescence analysis of Mouse lung tissue. 1, Smad2 Polyclonal Antibody (red) was diluted at 1:200 (4°C, overnight). 2, Cy3 labled secondary antibody was diluted at 1:300 (room temperature, 50min). 3, Picture B: DAPI (blue) 10min. Picture A: Target. Picture B: DAPI. Picture C: merge of A+B.

Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

Not for further distribution without written consent.Copyright © Abbkine, Inc.