



## Epac Polyclonal Antibody

Cat #: ABP53470

Size: 30µl /100µl /200µl

### Product Information

	<b>Product Name:</b> Epac Polyclonal Antibody		
	<b>Applications:</b> WB, ELISA		<b>Isotype:</b> Rabbit IgG
	<b>Reactivity:</b> Human, Mouse, Rat		
<b>REF</b>	<b>Catalog Number:</b> ABP53470	<b>LOT</b>	<b>Lot Number:</b> Refer to product label
	<b>Formulation:</b> Liquid		<b>Concentration:</b> 1 mg/ml
	<b>Storage:</b> Store at -20°C. Avoid repeated freeze / thaw cycles.		<b>Note:</b> Contain sodium azide.

**Background:** RAPGEF3 (Rap Guanine Nucleotide Exchange Factor 3) is a Protein Coding gene. Among its related pathways are Platelet activation, signaling and aggregation and Signaling by GPCR. GO annotations related to this gene include guanyl-nucleotide exchange factor activity and cAMP-dependent protein kinase regulator activity. An important paralog of this gene is RASGEF1A. uanine nucleotide exchange factor (GEF) for RAP1A and RAP2A small GTPases that is activated by binding cAMP. Through simultaneous binding of PDE3B to RAPGEF3 and PIK3R6 is assembled in a signaling complex in which it activates the PI3K gamma complex and which is involved in angiogenesis. Plays a role in the modulation of the cAMP-induced dynamic control of endothelial barrier function through a pathway that is independent on Rho-mediated signaling. Required for the actin rearrangement at cell-cell junctions, such as stress fibers and junctional actin. TPases are a group of enzymes that catalyze hydrolysis of the gamma phosphate bond in guanine triphosphate (GTP) to form guanine diphosphate (GDP). Mg<sup>2+</sup> ions are essential for catalytic activity. GTPases, often coupled to G proteins, are essential in signal transduction.

**Application Notes:** Optimal working dilutions should be determined experimentally by the investigator. Suggested starting dilutions are as follows: WB (1:500-1:2000), ELISA (1:40000). Not yet tested in other applications.

**Storage Buffer:** PBS containing 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.

**Storage Instructions:** Stable for one year at -20°C from date of shipment. For maximum recovery of product, centrifuge the original vial after thawing and prior to removing the cap. Aliquot to avoid repeated freezing and thawing.

**Note:** The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

