

CheKine™ Micro Fructose 1,6-Bisphosphate Aldolase (FBA) Activity Assay Kit

Cat #: KTB1332

Size: 48 T/48 S 96 T/96 S

	Micro Fructose 1,6-Bisphosphate Aldolase (FBA) Activity Assay Kit		
REF	Cat #: KTB1332	LOT	Lot #: Refer to product label
	Applicable sample: Animal and Plant tissues, Cells, Fungus, Plasma, Serum or other Liquid samples		
	Storage: Stored at 4°C for 12 months		

Assay Principle

Fructose 1,6-diphosphate aldolase (FBA)(EC.4.1.2.13) is an important enzyme involved in calvin cycle in glycolysis, gluconeogenesis, pentose phosphate pathway and photosynthesis, which catalyzes the reversible cleavage of fructose 1,6-diphosphate into dihydroxyacetone phosphate and glyceraldehyde 3-phosphate, and is widely used. CheKine™ Micro Fructose 1,6-Bisphosphate Aldolase (FBA) Activity Assay Kit can detect animal and plant tissues, cells, fungus, plasma, serum or other liquid samples. In this kit, FBA catalyzes fructose-1,6-diphosphate to produce glyceraldehyde-3-phosphate and dihydroxyacetone-3-phosphate, and NADH and dihydroxyacetone-3-phosphate are catalyzed to produce NAD and α-glycerophosphate under the action of triose phosphate isomerase and α-glycerophosphate dehydrogenase. The change of absorbance at 340 nm can reflect the activity of FBA.

Materials Supplied and Storage Conditions

Kit components	Size		Storage conditions
	48 T	96 T	
Extraction Buffer	60 mL	120 mL	4°C
Reagent I	6 mL	12 mL	4°C
Reagent II	3 mL	6 mL	4°C

Note: Before formal testing, it is recommended to select 2-3 samples with large expected differences for pre-experiment.

Materials Required but Not Supplied

- Microplate reader or ultraviolet spectrophotometer capable of measuring absorbance at 340 nm
- 96-well UV microplate or micro quartz cuvette, precision pipettes, disposable pipette tips, 1.5 mL EP tube
- Water bath, cryogenic centrifuge
- Deionized water
- Mortar or homogenizer (for tissue samples)

Reagent Preparation

Extraction Buffer: Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

Reagent I: Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

Reagent II: Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

Sample Preparation

Note: We recommend that you use fresh samples. If not assayed immediately, samples can be stored at -80°C for one month. When measuring, the temperature and time of thawing should be controlled. When thawing at room temperature, the sample should be thawed within 4 h.

1. Tissues: weigh about 0.1 g sample, add 1ml of Extraction Buffer, homogenize in ice bath, then centrifuge at 4°C for 10 min at 13,000 g, and take the supernatant and put it on ice for testing.
2. Cells or Fungus: Collect 5×10^6 cells or fungus into the centrifuge tube, wash cells or fungus with cold PBS, discard the supernatant after centrifugation; add 1 mL Extraction Buffer to ultrasonically disrupt the cells or fungus 5 min (power 20% or 200 W, ultrasonic 3 s, interval 7 s, repeat 30 times). Centrifuge at 13,000 g for 10 min at 4°C. Use supernatant for assay, and place it on ice to be tested.
3. Serum, Plasma or other Liquid samples: Test directly.

Note: If the protein concentration of the sample is need to determined, it is recommended to use Abbkine Cat #: KTD3001 Protein Quantification Kit (BCA Assay) to measure the protein concentration of the sample.

Assay Procedure

1. Preheat the microplate reader or ultraviolet spectrophotometer for more than 30 min, and adjust the wavelength to 340 nm, ultraviolet spectrophotometer was returned to zero with deionized water.
2. Working Reagent: each well need 100 μ L Reagent I and 50 μ L Reagent II, mix thoroughly before test.
3. Operation table (The following operations are operated in the 96-well UV microplate or micro quartz cuvette):

Reagent	Blank Well (μ L)	Test Well (μ L)
Deionized Water	20	0
Sample	0	20
Working Reagent	150	150

4. Mix thoroughly, measure the absorbance value A_1 at 3 min at 340 nm, and the absorbance value A_2 at 5 min at 37°C. The Test Well is marked as A_{Test} , the Blank Well is marked as A_{Blank} . Finally calculate $\Delta A = (A_{1\text{Test}} - A_{2\text{Test}}) - (A_{1\text{Blank}} - A_{2\text{Blank}})$.

Note: In order to guarantee the accuracy of experimental results, need to do a pre-experiment with 2-3 samples. If $A_{1\text{Test}}$ is greater than 1.0 with ΔA is less than 0.005, increase the sample quantity appropriately. If $A_{1\text{Test}}$ is less than 0.5, the sample can be appropriately diluted with Extraction Buffer, the calculated result multiplied by the dilution factor, or decrease the sample quantity appropriately. If ΔA is negative, the sample does not contain FBA or is degraded.

Data Analysis

Note: We provide you with calculation formulae, including the derivation process and final formula. The two are exactly equal. It is suggested that the concise calculation formula in bold is final formula.

Calculation of FBA activity:

A. 96-well UV plates calculation formula as below

(1) Calculated by protein concentration

Active unit definition: The consumption of 1 nmol of NADH per milligram of protein per min was defined as one unit of enzyme

activity.

$$\text{FBA (U/mg prot)} = [\Delta A \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (V_{\text{Sample}} \times \text{Cpr}) \div T = 1366.56 \times \Delta A \div \text{Cpr}$$

(2) Calculated by fresh weight of samples

Active unit definition: The consumption of 1 nmol of NADH per gram tissue per min was defined as one unit of enzyme activity.

$$\text{FBA (U/g fresh weight)} = [\Delta A \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (W \times V_{\text{Sample}} \div V_{\text{Total sample}}) \div T = 1366.56 \times \Delta A \div W$$

(3) Calculated by cell or fungus number

Active unit definition: The consumption of 1 nmol of NADH per 10^4 cell or fungus min was defined as one unit of enzyme activity.

$$\text{FBA (U/10}^4\text{)} = [\Delta A \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (n \times V_{\text{Sample}} \div V_{\text{Total sample}}) \div T = 1366.56 \times \Delta A \div n$$

(4) Calculated by volume of samples

Active unit definition: The consumption of 1 nmol of NADH per mL liquid per min was defined as one unit of enzyme activity.

$$\text{FBA (U/mL)} = [\Delta A \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (V_{\text{Sample}} \div V_{\text{Total sample}}) \div T = 1366.56 \times \Delta A$$

V_{Total} : total reaction volume, 1.7×10^{-4} L; ϵ : NADPH molar extinction coefficient, 6.22×10^3 L/mol /cm; d : the light path of the 96-well plate, 0.5 cm; V_{Sample} : sample volume added, 0.02 mL; $V_{\text{Total sample}}$: Extraction Buffer volume added, 1 mL; T : reaction time, 2 min; Cpr : sample protein concentration, mg/mL; W : weight of sample, g; n : Total number of cells or fungus, calculated in units of ten thousand.

B. Micro quartz cuvette calculation formula

The optical diameter d : 0.5 cm in the above calculation formula can be adjusted to d : 1 cm for calculation.

Typical Data

Weigh 0.105 g rabbit liver, follow the procedure, and perform the assay using a 96-well UV plate.

$$\Delta A = (A_{1\text{Test}} - A_{2\text{Test}}) - (A_{1\text{Blank}} - A_{2\text{Blank}}) = (1.534 - 1.361) - (1.526 - 1.523) = 0.170, \text{ FBA (U/g)} = 1366.56 \times \Delta A \div W = 2212.5 \text{ U/g.}$$

Recommended Products

Catalog No.	Product Name
KTB3030	CheKine™ Micro Alcohol Dehydrogenase (ADH) Activity Assay Kit
KTB1560	CheKine™ Micro Alcohol Acyltransferase (AAT) Activity Assay Kit
KTB1270	CheKine™ Micro Pyruvate Dehydrogenase (PDH) Activity Assay Kit

Disclaimer

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes. For your safety and health, please wear a lab coat and disposable gloves.