



## CheKine™ Micro Mitochondrial Complex IV Activity Assay Kit

Cat #: KTB1880

Size: 48 T/96 T

	<b>Micro Mitochondrial Complex IV Activity Assay Kit</b>		
<b>REF</b>	<b>Cat #:</b> KTB1880	<b>LOT</b>	<b>Lot #:</b> Refer to product label
	<b>Applicable samples:</b> Animal and Plant Tissues, Cells		
	<b>Storage:</b> Stored at -20°C for 6 months, protected from light		

### Assay Principle

Mitochondrial respiratory chain complex IV, also known as cytochrome C oxidase, is also a common component of the main and branch pathways of the mitochondrial respiratory chain. It is responsible for catalyzing the oxidation of reduced cytochrome C, and ultimately transferring electrons to oxygen to produce water. CheKine™ Micro Mitochondrial Complex IV Activity Assay Kit provides a convenient tool for detection of Mitochondrial complex IV Activity. The principle is Reduced cytochrome C has characteristic light absorption at 550 nm. Mitochondrial respiratory chain complex IV could catalyze reduced cytochrome C to oxidized cytochrome C. Therefore, the decreasing rate of light absorption at 550 nm can reflect the enzyme activity of mitochondrial respiratory chain complex IV. It can be used to determine animal, plant tissues and cell samples.

### Materials Supplied and Storage Conditions

Kit components	Size		Storage conditions
	48 T	96 T	
Reagent I	50 mL	100 mL	4°C
Reagent II	10 mL	20 mL	4°C
Reagent III	1 mL	2 mL	4°C, protected from light
Reagent IV	10 mL	20 mL	4°C
Reagent V	1	1	-20°C, protected from light
Reagent VI	1	1	-20°C, protected from light

### Materials Required but Not Supplied

- Microplate reader or visible spectrophotometer capable of measuring absorbance at 550 nm
- Incubator, ice maker, refrigerated centrifuge
- 96-well plate or microglass cuvette, precision pipettes, disposable pipette tips
- Deionized water

- Homogenizer (for tissue samples)

## Reagent Preparation

**Reagent I** : Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

**Reagent II** : Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C.

**Reagent III** : Ready to use as supplied. Equilibrate to room temperature before use. Store at 4°C, protected from light.

**Working Solution**: Transfer the Reagent V and Reagent VI respectively to the tube of Reagent IV, and mixing well before use.

Then incubated the mixture at 37°C for 5 min if the detected samples are from mammalian, or incubated at 25°C for 5 min if the samples are from another species.

## Sample Preparation

**Note: Fresh samples are recommended to ensure enzyme activity.**

Extraction of Mitochondrial Respiratory Chain Complex IV:

1. Accurately weigh 0.1 g tissue or collect  $5 \times 10^6$  cells, add 1 mL Reagent I and 10  $\mu$ L Reagent III, homogenize on ice.
2. Centrifuge the homogenate with 600 g for 5 min at 4°C, collect the supernatant to a new centrifuge tube and discard the pellet.
3. Centrifuge the supernatant again with 11,000 g for 10 min at 4°C. The pellet is the extracted mitochondria, which could be used to do step 5.
4. (Optional) The supernatant is cytoplasmic extract, which can be used as sample to determine mitochondrial respiratory chain complex IV leaking from mitochondria to judge the effect of mitochondrial extraction.
5. Add 200  $\mu$ L Reagent II and 2  $\mu$ L Reagent III to the pellet, resuspend the pellet sufficiently, and use it to detect the activity of mitochondrial respiratory chain complex IV in the next step.

## Assay Procedure

1. Preheated the microplate reader or visible spectrophotometer for more than 30 min, and adjust the wavelength to 550 nm, visible spectrophotometer was returned to zero with deionized water.
2. Successively add 200  $\mu$ L Working Solution and 10  $\mu$ L sample into the 96-well plate or microglass cuvette, mix well. Immediately read the initial absorbance value (0 min) at 550 nm as  $A_1$ , and then read again after 1 min as  $A_2$ . Finally calculate  $\Delta A = A_1 - A_2$ .

**Note: In order to guarantee the accuracy of experimental results, need to do a pre-experiment with 1-2 samples. If  $\Delta A$  is too high (above 1.0), the samples should be dilute with Reagent II and then measured again. Pay attention to multiply by the dilution factor when calculating the result. If  $\Delta A$  is too small, the sensitivity can be improved by increasing the sample volume added. If  $\Delta A$  is negative, it means that complex IV is not contained in the sample or has been degraded.**

## Data Analysis

**Note: We provide you with calculation formulae, including the derivation process and final formula. The two are exactly equal. It is suggested that the concise calculation formula in bold is final formula.**

1. 96-well plates calculation formula as below

(1) Calculated by fresh weight of samples

Unit definition: One enzyme activity unit defines as the consumption of 1 nmol reduced cytochrome C in 1 g tissue per min in the reaction system.

Calculate the activity of Complex IV in the supernatant:

The activity of supernatant Complex IV (U/g fresh weight) =  $[\Delta A_1 \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (W \div V_{\text{Extraction}} \times V_{\text{Sample}}) \div T = \mathbf{2,221 \times \Delta A_1 \div W}$

Calculate the activity of Complex IV of the mitochondrial pellet:

The activity of pellet Complex IV (U/g fresh weight) =  $[\Delta A_2 \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (W \div V_{\text{Resuspended}} \times V_{\text{Sample}}) \div T = \mathbf{444 \times \Delta A_2 \div W}$

Calculate the total activity of Complex IV in sample:

The total activity of Complex IV in sample is the sum of the supernatant and pellet.

Total activity (U/g fresh weight)= $2,221 \times \Delta A_1 \div W + 444 \times \Delta A_2 \div W$

(2) Calculated by cell density

Unit definition: Every 10,000 cells consume 1 nmol reduced cytochrome C per minute that is defined as one enzyme activity.

The activity of Complex IV (U/10<sup>4</sup> cells)= $[\Delta A \times V_{\text{Total}} \div (\epsilon \times d) \times 10^9] \div (V_{\text{Sample}} \div V_{\text{Resuspend}} \times 500) \div T = 0.888 \times \Delta A$

Where:  $V_{\text{Total}}$ : total reaction volume, 2.1×10<sup>-4</sup> L;  $\epsilon$ : reduced cytochrome C molar extinction coefficient, 19.1×10<sup>3</sup> mol/L/cm; d: 96-well plate diameter, 0.5 cm; 10<sup>9</sup>: Unit conversion factor, 1 mol=10<sup>9</sup> nmol;  $V_{\text{Sample}}$ : sample volume added, 0.01 mL; T: reaction time, 1 min;  $\Delta A_1$ : determination value of the supernatant; W: sample weight, g;  $V_{\text{Extraction}}$ : sample extract volume, 1.01 mL;  $\Delta A_2$ : determination value of pellet;  $V_{\text{Resuspended}}$ : volume of the resuspend pellet 0.202 mL; 500: total number of cells, 5×10<sup>6</sup>.

2. Microglass cuvette calculation formula

The optical diameter d: 0.5 cm in the above calculation formula can be adjusted to d: 1 cm for calculation

## Typical Data

Examples:

1. Test 0.1g mouse brain tissue, prepared the sample following the above protocol and measured with the 96-well microplate:

$\Delta A_1 = A_1 - A_2 = 0.4568 - 0.4446 = 0.0122$ ,  $\Delta A_2 = A_3 - A_4 = 0.4373 - 0.4188 = 0.0185$

2. Calculated by fresh weight of samples,

Complex IV activity of the supernatant (U/g fresh weight)= $2,221 \times \Delta A_1 \div W = 2,221 \times 0.0122 \div 0.1 = 270.962$  U/g

Complex IV activity of the pellet (U/g fresh weight) = $444 \times \Delta A_2 \div W = 444 \times 0.0185 \div 0.1 = 82.14$  U/g

The total Complex IV activity (U/g fresh weight)= $2,221 \times \Delta A_1 \div W + 444 \times \Delta A_2 \div W = 270.962 + 82.14 = 353.102$  U/g

## Recommended Products

Catalog No.	Product Name
KTB1850	CheKine™ Micro Mitochondrial Complex I Activity Assay Kit
KTB1860	CheKine™ Micro Mitochondrial Complex II Activity Assay Kit
KTB1870	CheKine™ Micro Mitochondrial Complex III Activity Assay Kit
KTB1890	CheKine™ Micro Mitochondrial Complex V Activity Assay Kit

## Disclaimer

The reagent is only used in the field of scientific research, not suitable for clinical diagnosis or other purposes.