



Technical support: support@abbkine.com

Website: https://www.abbkine.com

Recombinant Human IL-8 protein

Cat #:PRP1022 Size: 10µg /50µg/100µg /1mg

Product Information

	Product Name: Recombinant Human IL-8 protein		
	Reconstitution: Use the buffer provided to reconstitute the lyophilized Human IL-8 protein		
REF	Catalog Number: PRP1022	LOT	Lot Number: Refer to product label
	Purity: ≥ 98 % as determined by SDS-PAGE		
Ŷ	Storage: Store at -20°C		Preparation method: E.coli
	Shipping: The product is shipped at ambient temperature.		

Background: Interleukin-8 (IL-8), also known as CXCL8, GCP-1, and NAP-1, belongs to the neutrophil-specific CXC family of chemokines. This chemokine is secreted by a variety of cell types including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, and various tumor cell lines in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation.

Sequence: Amino acid sequence derived from Human IL-8 isoform (P10145)(Ser28-Ser99) was expressed.

<u>Protein length</u>: The recombinant Human IL-8 consists of 72 amino acids (N-Met) and has a predicted molecular mass of 8 kDa. The apparent molecular mass of the Human IL-8 protein is approximately 8 kDa in SDS-PAGE under reducing conditions.

Biological Activity: Testing in progress.

Endotoxin: < 1 EU per μg of the protein as determined by the LAL method.


Formulation: Lyophilized from sterile 20 mM Tris, 200 mM NaCl pH 7.0.

Storage Instructions: Lyophilized Human IL-8 protein product should be stored desiccated below -18°C for 12 months. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and below -18°C for 3 months. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). aliquot and store below -18°C. Please prevent freeze-thaw cycles.

Usage notes: Always centrifuge tubes before opening. It is recommended to centrifuge at 12,000 rpm for 1 minute. Then



reconstitute the lyophilized Human IL-8 protein with the buffer provided not less than 100µg/ml, which can then be further diluted with other aqueous solutions.



<u>Note:</u> The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

