



Technical support: support@abbkine.com

Website: https://www.abbkine.com

Human BMP-13 Protein, His tag (Animal-Free)

Cat #: PRP1205 Size: 5 μg/20 μg/100 μg/1 mg

Product Information

	Product Name: Human BMP-13 Protein, His tag (Animal-Free)		
REF	Catalog Number: PRP1205	LOT	Lot Number: Refer to product label
	Purity: > 98 % as determined by SDS-PAGE		
Å	Storage: Store at -20°C		Preparation method: E.coli
	Shipping: The product is shipped at ambient temperature.		

Background: Bone Morphogenetic Protein-13 (BMP-13), known as Growth Differentiation Factor 6 (GDF6), is an extracellular multifunctional cytokine that is also a member of the TGFβ family. BMP-13 can bind with the TGFβ receptor and induce SMAD protein signal transduction. BMP-13 is a regulatory protein that can affect the growth and differentiation of developing embryos. Moreover, it plays a role in regulating the patterning of the ectoderm and controlling eye development.

<u>Sequence:</u> Amino acid sequence derived from Human BMP-13 (Met335-Arg455) (Q6KF10) was expressed with 6×His tag at the C-terminus.

Protein length: The recombinant Human BMP-13 consists of 121 amino acids and predicts a molecular mass of 14.5 kDa.

<u>Biological Activity</u>: Measured by its ability to induce alkaline phosphatase production by ATDC5 cells. The ED₅₀ for this effect is 63-240 ng/mL.


Endotoxin: <0.1 EU per 1 µg of the protein by the LAL method.

Formulation: Lyophilized from 20 mM sodium citrate, 0.2 M NaCl, pH 3.5.

Storage Instructions: Lyophilized protein product should be stored desiccated below -18°C. Upon reconstitution, the protein should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (5% HSA, 10%FBS or 0.1%BSA). Please prevent freeze-thaw cycles.

<u>Usage notes</u>: Always centrifuge tubes before opening. It is recommended to reconstitute the lyophilized recombinant protein to a concentration of 0.1-1 mg/mL using the buffer we provided, and keep at room temperature for at least 20 min to fully dissolve. Please avoid vortex vigorously.





Note: The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.

