





## Terminal Deoxynucleotidyl Transferase (TdT)

Cat #: PRP3002

Size: 40 µg/ 100 µg

### Product Information

	<b>Product Name:</b> Terminal Deoxynucleotidyl Transferase (TdT)		
	<b>Catalog Number:</b> PRP3002		<b>Lot Number:</b> Refer to product label
	<b>Purity:</b> > 90% as determined by SDS-PAGE		
	<b>Storage:</b> Store at -20°C		<b>Expressed Host:</b> E. coli
	<b>Shipping:</b> Gel pack with blue ice		

**Background:** Terminal deoxynucleotidyl transferase (TdT), also known as terminal transferase, is a specialized DNA polymerase expressed in immature, pre-B, pre-T lymphoid cells, and acute lymphoblastic leukemia/lymphoma cells. Generally, TdT catalyses the addition of nucleotides to the 3' terminus of a DNA molecule. Unlike most DNA polymerases, it does not require a template. The preferred substrate of this enzyme is a 3'-overhang, but it can also add nucleotides to blunt or recessed 3' ends.

**Sequence:** Amino acid sequence derived from Bovine TdT (X04122) (Mer1-Ala520) was expressed.

**Protein length:** The recombinant Bovine TdT consists of 520 amino acids and migrates with an apparent molecular mass of 58.3 kDa as estimated in SDS-PAGE under reducing conditions.

**Unit Definition:** One unit of the enzyme catalyzes the incorporation of 1 nmol of deoxythymidylate into a polynucleotide fraction in 60 min at 37°C.

**Storage buffer:** Liquid in sterile 50 mM K<sub>2</sub>HPO<sub>4</sub>, 100 mM NaCl, 1 mM DTT, 0.1% Tween20, 1% BSA, 50% Glycerol, pH 6.5.


#### Formulation:

Components	Size 1	Size 2
Terminal Deoxynucleotidyl Transferase (TdT), 1 mg/mL	40 µg	100 µg
Equilibration Buffer (5×)	0.5 mL	1 mL

**Equilibration Buffer (5×):** 1 M potassium cacodylate, 125 mM Tris, 1 mM DTT, 5 mM CoCl<sub>2</sub>.

**Usage notes:** The Equilibration Buffer (5×) stock solution is stored at low temperature, resulting in a small amount of component precipitation. Please invert and mix before use. And Equilibration Buffer (5×) contains cacodylate and cobalt chloride, highly toxic chemicals. After contact with skin, wash immediately with plenty of water. In case of accident or if you feel unwell, seek medical advice immediately.

Reference system of TUNEL:	Equilibration Buffer (5×)	10 μL
	Fluorescence-labeled dUTP	5 μL
	ddH <sub>2</sub> O	33 μL
	Terminal Deoxynucleotidyl Transferase (TdT)	2 μL



**Note:** The product listed herein is for research use only and is not intended for use in human or clinical diagnosis. Suggested applications of our products are not recommendations to use our products in violation of any patent or as a license. We cannot be responsible for patent infringements or other violations that may occur with the use of this product.